SUBJECT	DATE	PAGE	ITEM NO.
Sample Collection: Requirements for Drinking Water	11/18/19	1 of 9	241

# ANALYTE CONTAINER PRESERVATION MAXIMUM HOLDING TIME

Note 1: Consumer collected samples may be left unpreserved for up to 14 days.

Note 2: ELAP offers Nitrate or Nitrite only for accreditation. ELAP does not offer combined Nitrate-Nitrite. The preservation and holding time requirements for combined Nitrate-Nitrite is Cool to 4°C, H<sub>2</sub>SO<sub>4</sub> to pH<2, and 28 days.

Note 3: Per NELAC/TNI standards, "All samples, which require thermal preservation, shall be considered acceptable if the arrival temperature is either within  $\pm$  2 °C of the required temperature or the method specified range. For a sample with a specified temperature of 4 °C, sample with a temperature ranging from just above the freezing temperature of water to 6 °C shall be acceptable..."

#### **Inorganic Tests:**

Alkalinity	P,G	Separate bottle completely filled to the exclusion of air. Cool, 4°C	14 days
Antimony	P,G	HNO₃ to pH<2	6 months
Arsenic	P,G	HNO₃ to pH<2	6 months
Barium	P,G	HNO₃ to pH<2	6 months
Beryllium	P,G	HNO₃ to pH<2	6 months
Inorganic Tests:			
Bromate	P,G	50 mg IL EDA	28 days
Cadmium	P,G	HNO <sub>3</sub> to pH<2	6 months
Calcium	P,G	HNO₃ to pH<2	6 months
Chloride	P,G	None	28 days
Chlorine Residual	P,G	None	immediately
Chlorite	P,G	50 mg IL EDA, Cool, 4 <sup>0</sup> C	14 days
Chromium	P,G	HNO <sub>3</sub> to pH<2	6 months
Color	P,G	Cool, 4 <sup>o</sup> C	48 hours

SUBJECT	DATE	PAGE	ITEM NO.
Sample Collection: Requirements for Drinking Water	11/18/19	2 of 9	241

<u>ANALYTE</u>	CONTAINER	PRESERVATION	MAXIMUM HOLDING TIME
Conductivity	P,G	Cool, 4 <sup>o</sup> C	28 days
Copper	P,G	HNO <sub>3</sub> to pH<2 Note 1	6 months
Cyanide	P,G	Cool, 4 <sup>o</sup> C, NaOH to pH=12, 1.2 g/L ascorbic acid	14 days
Fluoride	P,G	None	28 days
Lead	P,G	HNO $_3$ to pH<2 $^{Note\ 1}$	6 months
Mercury	P,G	HNO <sub>3</sub> to pH<2	28 days
Nickel	P,G	HNO <sub>3</sub> to pH<2	6 months
Nitrate Note 2	P,G	Cool, 4 <sup>o</sup> C	48 hours
Chlorinated Samples only	P,G	Cool, 4 <sup>o</sup> C	14 days
Nitrite	P,G	Cool, 4 <sup>o</sup> C	48 hours
рН	P,G	None	immediately
Phosphorus (as Orthophosphate)	P,G	Cool, 4 <sup>o</sup> C	48 hours
Selenium	P,G	HNO₃ to pH<2	6 months
Silica	Р	Cool, 4 <sup>o</sup> C	28 days
Silver	P,G	HNO₃ to pH<2	6 months
Sodium	P,G	HNO₃ to pH<2	6 months
Sulfate	P,G	Cool, 4 <sup>o</sup> C	28 days
Thallium	P,G	HNO₃ to pH<2	6 months
Total Filterable Residue	P,G	Cool, 4 <sup>o</sup> C	7 days
Turbidity	P,G	Cool, 4 <sup>o</sup> C	48 hours

SUBJECT	DATE	PAGE	ITEM NO.
Sample Collection: Requirements for Drinking Water	11/18/19	3 of 9	241

Cample Collection: Hequirements to	Brilliang Water	11/10/13	271
<u>ANALYTE</u>	CONTAINER	PRESERVATION	MAXIMUM HOLDING TIME
UV <sub>254</sub> Absorbance	P,G	Cool, 4°C	48 hours
Organic Tests:			
Trihalomethanes  Bromodichloromethane Bromoform Chlorodibromomethane Chloroform	Glass with Teflon-lined Septum	0.008% Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> , Cool, 4 <sup>o</sup> C	14 days
Volatile Halocarbon and Volatile Aromatics:  Methy-tert-butyl ether	Glass with Teflon-lined Septum	Ascorbic Acid (25 mg/40 ml) added to empty sample bottle then add 1:1 HCl to pH<2. Cool, 4 <sup>o</sup> C	14 days
Microextractables: Method 504.1	Glass with Teflon-lined Septum	Cool, 4 <sup>o</sup> C, 3 mg Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> per 40 ml vial	14 days
Alachlor Aldrin Atrazine Chlordane Dieldrin Heptachlor Heptachlor epoxide Hexachlorobenzene Hexachlorocyclopentadiene Lindane Methoxychlor Metolachlor Metribuzin PCB's Simazine Toxaphene	40-ml glass vial with cap liner	3 mg Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> , Cool, 4 <sup>o</sup> C	7 days
Method 506 analytes Di-(2-ethylhexyl)adipate Di-(2-ethylhexyl) phthalate	1-L (or qt.) amber glass with TFE lined cap	60 mg Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> , Cool, 4 <sup>o</sup> C	14 days until extraction, then 14 days after extraction

SUBJECT	DATE	PAGE	ITEM NO.
Sample Collection: Requirements for Drinking Water	11/18/19	4 of 9	241

· · · · · · · · · · · · · · · · · · ·		•	•
<u>ANALYTE</u>	CONTAINER	PRESERVATION	MAXIMUM HOLDING TIME
Method 507 analytes  Alachlor Atrazine Butachlor Chlordane Metolachlor Metribuzine Propachlor Simazine	1-L Borosilicate glass, graduated, with TFE lined cap	80 mg Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> , Cool, 4 <sup>o</sup> C, Protect from light after extraction	14 days until extraction, then 14 days
Aldrin Chlordane Dieldrin Endrin Heptachlor Heptachlor epoxide Hexachlorobenzene Hexachlorocyclopentadiene Lindane Methoxychlor Metribuzin PCB's Toxaphene	1-L Borosilicate glass, graduated, with TFE lined cap	80 mg Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> , Cool, 4 <sup>o</sup> C, Protect from light	7 days until extraction, then 14 days after extraction
Method 508A PCB's, Total as decachlorobiphenyl	1-L glass, with TFE lined cap	Cool, 4 <sup>o</sup> C	14 days until extraction, then 30 days after extraction
Method 508.1 All	1-L glass with TFE lined cap	50 mg Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> then 1:1 HCl to pH<2, Cool, 4 <sup>o</sup> C	14 days until extraction then 30 days after extraction
Method 1613 2,3,7,8-TCDD	1-L amber glass with TFE lined cap	80 mg Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> , Cool, 4 <sup>o</sup> C, Protect from light pH 7-9	one year until extraction, then one year after extraction

SUBJECT	DATE	PAGE	ITEM NO.
Sample Collection: Requirements for Drinking Water	11/18/19	5 of 9	241

Sample Collection: Requirements to	i Diliking water		241
ANALYTE	CONTAINER	PRESERVATION	MAXIMUM HOLDING TIME
Method 515.1: 515.2, 515.3 Chlorinated Acids	1-L Borosilicate glass, graduated, with TFE lined cap	80 mg Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> , Cool, 4 <sup>o</sup> C Protect from light	14 days until extraction, then 14 days after extraction
Method 524.3	40-ml VOA vial	See section 8.1 and 8.4 of method.	Analyze within 14 days of collection
Alachlor Aldrin Atrazine Benzo(a)pyrene Butachlor Chlordane (Technical) Dieldrin Di(2ethylhexyl)adipate Di(2-ethylhexyl) phthalate Endrin Heptachlor Heptachlor Epoxide Hexachlorobenzene Hexachlorocyclopentadiene Lindane Methoxychlor Metolachlor Metribuzin Pentachlorophenol Propachlor Simazine Toxaphene	Refrigerated glass sample containers - Sampling must be free of plastic tubing, gaskets, etc. that may leach analytes into water.	Cool, 4 C, Remove CI residual; adjust pH<2 with6 N HCI	Extract within14 days. Analyze within 30 days of sample extraction
Method 531.1  Methylcarbamate pesticides	60-ml vial with PTFE silicone faced septa	1.8 ml acetic acid buffer, 4.8 mg Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> , pH=3±0.2 for 4 analytes, Ship at 4 <sup>o</sup> C, Store at -	28 days

10<sup>0</sup>C

SUBJECT	DATE	PAGE	ITEM NO.
Sample Collection: Requirements for Drinking Water	11/18/19	6 of 9	241

ANALYTE	CONTAINER	PRESERVATION	MAXIMUM HOLDING TIME
Method 531.2  Methylcarbamate pesticides	60-ml vial with PTFE silicone faced septa	Use Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> & $C_6H_7KO_7$ to pH~3.8 for 4 analytes, Ship at ≤10°C, Store in dark at ≤6°C	28 days
Glyphosate	60-ml vial PTFE faced Silicone	6 mg Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> , Cool, 4 <sup>o</sup> C, Protect from light	14 days
Endothall	40-ml amber glass vial with TFE lined cap	Cool, 4 <sup>o</sup> C, Protect from light	7 days
Diquat	1-L amber plastic or silanized glass with screw cap	100 mg Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> H <sub>2</sub> SO <sub>4</sub> to pH=2, Cool, 4 <sup>o</sup> C, Protect from light	7 days until extraction, then 21 days after extraction
Benzo(a)pyrene	1-L (or qt.) amber glass with TFE lined cap	100 mg Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> , 1:1 HCl to pH<2, Cool, 4 <sup>o</sup> C; Protect from light	7 days until extraction then 30 (40 for Method550.1) days after extraction

SUBJECT	DATE	PAGE	ITEM NO.
Sample Collection: Requirements for Drinking Water	11/18/19	7 of 9	241

Sample Collection: Requirements for	Drinking Water	11/18/19 7 of 9	241
<u>ANALYTE</u>	CONTAINER	PRESERVATION	MAXIMUM HOLDING TIME
Alachlor Atrazine Bromochloromethane Bromodichloromethane Bromoform Carbon Tetrachloride Chloroform Dibromochloromethane 1,2-Dibromo-3- chloropropane [DBCP] 1,2 Dibromoethane [EDB] Endrin Heptachlor Heptachlor epoxide Hexachlorobenzene Hexachlorocyclopentadiene Lindane Methoxychlor Metalochlor Metribuzin Simazine Tetrachloroethylene 1,1,1-Trichloroethane 1,1,2-Trichloroethane Trichloroethylene	60 ml glass vials with Teflon lined Septum	Sodium Sulfite or Ammonium Chloride (for microextractables), pH 4.5-5.5 with phosphate buffer, Cool, 4°C	14 days until extraction, then 14 days after extraction
Method 552.1  Dalapon  Monochloroacetic acid Dichloroacetic acid Trichloroacetic acid Monobromoacetic acid	Amber glass with TFE liner	Add NH <sub>4</sub> Cl to a concentration of 100 mg/L in sample, Cool, 4 <sup>o</sup> C	Extract within 28 days of collection. Analyze extract within 48 hours if stored at 4°C or less

Dibromoacetic acid Bromochloroacetic Acid or less.

SUBJECT	<u>DATE</u>	PAGE	ITEM NO.
Sample Collection: Requirements for Drinking Water	11/18/19	8 of 9	241

ANALYTE	CONTAINER	PRESERVATION	MAXIMUM HOLDING TIME
Dalapon Monochloroacetic acid Dichloroacetic acid Trichloroacetic acid Monobromoacetic acid Dibromoacetic acid Bromochloroacetic Acid	Amber glass with TFE liner	Add NH <sub>4</sub> Cl to a concentration of 100 mg/L in sample, Cool, 4 <sup>o</sup> C	Extract within 14 days of collection. Analyze extract within 7 days if stored dark at 4°C or less or 14 days if - 10°C or less. See section 8.3 of method.
Method 552.3  Dalapon Monochloroacetic acid Dichloroacetic acid Trichloroacetic acid Monobromoacetic acid Dibromoacetic acid Bromochloroacetic Acid	Amber glass with TFE liner	Add NH <sub>4</sub> Cl to a concentration of 100 mg/L in sample, Cool, 4 <sup>o</sup> C. Extracts stored at -10 <sup>o</sup> C.	Extract within 14 days. See section 8.4 of method.
Method 555  2,4-D  Dicamba  Pichloram  2,4,5-TP	Glass with TFE liner	Acidify to pH2 with 1:1 HCl, Dechlorinate with 5 mg NaSO <sub>3</sub> per 100mL sample, Cool, 4 <sup>o</sup> C, Protect from light	Analyze after extraction, within 14 days of collection
Microscopical Tests:			
Asbestos	P,G	Cool, 4 <sup>o</sup> C	48 hours
Asbestos	P,G	Preserved with 10 gm/L of O <sub>3</sub> , and UV treatment	6 months

SUBJECT	DATE	PAGE	ITEM NO.
Sample Collection: Requirements for Drinking Water	11/18/19	9 of 9	241

ANALYTE CONTAINER PRESERVATION MAXIMUM HOLDING TIME

Instructions for containers, preservation procedures and holding times as specified in Method 100.2 must be adhered to for all compliance analyses including those conducted with Method 100.1.

#### **Radiological Tests:**

Gross Alpha	P,G	HCl or HNO₃ to pH<2	6 months (1yr for composites)
Gross Beta	P,G	HCl or HNO₃ to pH<2	6 months
Strontium-89	P,G	HCl or HNO₃ to pH<2	6 months
Strontium-90	P,G	HCl or HNO₃ to pH<2	6 months
Radium-226	P,G	HCl or HNO₃ to pH<2	6 months (1yr for composites)
Radium-228	P,G	HCl or HNO₃ to pH<2	6 months (1yr for composites)
Radon-222	Glass with teflon-lined septum	Cool, 4 <sup>o</sup> C**	3 days*
Radioactive Cesium	P,G	HCl to pH<2	6 months
lodine-131	P,G	None	7 days
Tritium	G	None	6 months
Uranium	P,G	HCl or HNO₃ pH<2	6 months (1yr for composites)
Photon Emitters	P,G	HCl or HNO₃ pH<2	6 months

<sup>\*</sup> Hold time varies based on the method used. If using Standard Methods, the hold time is 4 days. If using the EPA method, the hold time is 3 days.

<sup>\*\*</sup> Regardless of the method used, the samples are to be stored in a cooler or equivalent insulated container.